HALEON

Personalized Pain Management Solutions

Explore the range of Advil products, built on science

	Activity DUAL ACTION Deceminion Activity 25 ong add Austrin ISAND 25 on	Studied burden Epsiles 20179 Par Reiseer Feuer Reducer (NSAU)	buporten Tablets. 200 mg Pan Relever / Fever Reducer (NSAID) Contre Tablets	
	Advil Dual Action	Advil Liqui-Gels	Advil Coated Tablets	Advil Targeted Relief Pain Relieving Cream
Key Features	The 1st FDA-approved pain reliever to combine ibuprofen and acetaminophen Greater analgesia than when each medicine is taken alone	Rapid pain relief starting in minutes Available with an osteoarthritis cap	Trusted pain relief for 35+ years Available with an osteoarthritis cap	Starts working on contact Lasts up to 8 hours
Top Indications	Backache, dental pain, menstrual pain and other acute pains	Headache, musculoskeletal pain, fever, pain from common cold and minor pain of arthritis	Minor pain of arthritis, fever, headache, pain from common cold and musculoskeletal pain	Acute aches and pains including: -Back and body aches -Strains, sprains and sore muscles
Ingredients	Acetaminophen 250 mg Ibuprofen 125 mg (per tablet)	Ibuprofen 200 mg (per tablet)	Ibuprofen 200 mg (per tablet)	Menthol 10% Capsaicin 0.025% Camphor 3.1% Methyl Salicylate 15% Does not contain ibuprofen
Duration	8 hours	6-8 hours	6-8 hours	Up to 8 hours
Dosage	Adults and children 12 years and older: 2 tablets every 8 hours Maximum of 6 tablets over 24 hours	Adults and children over 12: 1 to 2 tablets every 4-6 hours as needed Maximum of 6 tablets over 24 hours	Adults and children over 12: 1 to 2 tablets every 4-6 hours as needed Maximum of 6 tablets over 24 hours	Adults and children over 12 Apply to affected area 3-4 times a day for up to 7 days

HALEON

Personalized Pain Management Solutions

Clinically proven pain relief for arthritis, headaches and migraine

	ARTHRITIS PAIN Voltaren Gel	Excedrin Extra Strength	Excedrin Migraine
Key Features	Penetrates deep through the skin at the site of pain ^{1,2} Improves mobility ^{2,3} Low systemic exposure ⁴ The American College of Rheumatology strongly recommends topical NSAIDs as a 1st-line treatment for osteoarthritis pain of the knee	Fast-acting formula – for some, headache relief starts in as little as 15 minutes	Provides effective relief of moderate-to-severe migraines with just 1 dose Starts relieving migraine pain in 30 minutes for some patients The #1 Neurologist recommended OTC*
Top Indications	Relieves arthritis pain in the hand, wrist, elbow, foot, ankle, and knee	Headaches	Migraines
Ingredients	1% diclofenac sodium	Acetaminophen 250 mg Aspirin 250 mg Caffeine 65 mg	Acetaminophen 250 mg Aspirin 250 mg Caffeine 65 mg
Duration	Approved for use up to 21 days After 21 days, patients should consult a physician	Up to 6 hours	Up to 24 hours
Dosage	Patients should use Voltaren 4 times a day, every day Do not use on more than 2 body areas at the same time	Adults and children 12 years and over: Take 2 tablets every 6 hours with a glass of water Maximum of 8 tablets over 24 hours	Adults 18+: Take 2 tablets per day with a glass of water Maximum of 2 tablets over 24 hours

*Among OTC medicines for migraines.

References: 1. Barthel HR, et al. Randomized controlled trial of diclofenac sodium gel in knee osteoarthritis. *Semin Arthritis Rheum*. 2009;39(3):203-212. **2.** Altman RD, et al. Diclofenac sodium gel in patients with primary hand osteoarthritis: a randomized, doubleblind, placebo-controlled trial. *J Rheumatol*. 2009;36(9):1991-1999. **3.** Zacher J, et al. Topical diclofenac and its role in pain and inflammation: an evidence-based review. *Curr Med Res Opin*. 2008;24:925-950. **4.** Riess W, et al. The percutaneous absorption of diclofenac. *Arzneimittelforschung*. 1986;36:1092-1096.

©2024 HALEON group of companies or its licensor. All rights reserved. PM-US-ADV-24-00175

Scan here to learn more

